

CLAIMS

1. Formulation (F) intended for use in an operation of rinsing (R) textile fiber articles (S) by means of an aqueous or aqueous-alcoholic medium (MR),
- 5 said formulation (F)
- comprising at least one active substance (A) comprising at least one solid or liquid organic or organosilicon material in particulate form and a vehicle (V) comprising at least one organic polymer,
- 10 capable of taking said active substance (A) to the surface of said textile fiber articles (S) in the rinsing operation (R),
- in the form:
 - of a stable dispersion, with a pH of from 2 to
- 15 5, of said active substance (A) in an aqueous or aqueous-alcoholic medium (MAV) comprising said vehicle (V), or
- in a solid form obtained by drying said dispersion,
- 20 the nature of the active substance (A), of the aqueous or aqueous-alcoholic medium (MAV), and of the vehicle (V) being such that
- * the active substance (A)
 - is insoluble in the medium (MAV),
- 25 • has an overall zero or cationic charge in the medium (MAV),

- is stabilized in the medium (MAV) by means of a cationic surfactant (TAC), it being possible for said cationic surfactant (TAC) to be wholly or partly replaced by a nonionic surfactant when the material constituting the active substance (A) is intrinsically cationic or intrinsically potentially cationic in the medium (MAV),
 - remains insoluble in the rinsing medium (MR);
- the vehicle (V)
- is soluble or dispersible in the medium (MAV) and in the rinsing medium (MR)
 - has an overall cationic or zero ionic charge in the medium (MAV),
 - at the pH of the rinsing operation in the rinsing medium (MR) is capable of developing anionic charges in sufficient quantity to destabilize the active substance (A) in the rinsing medium (MR).

2. Formulation according to claim 1), characterized in that the rinsing medium (MR) has a pH of from 5.5 to 8.

3. Formulation according to claim 1) or 2), characterized in that the material constituting the active substance (A) is an oil or a meltable solid.

4. Formulation according to any one of claims 1) to 3), characterized in that the particles of active substance (A) have an average diameter ranging from 10 nm to 200 μ m, preferably from 10 nm to 5 μ m and
5 more preferably from 10 nm to 2 000 nm.

5. Formulation according to any one of claims 1) to 4), characterized in that the active substance (A) comprises material having lubricating properties with regard to textile fiber articles.

10 6. Formulation according to any one of claims 1) to 5), characterized in that the material constituting the active substance (A) is a polyorganosiloxane selected from

- nonionic polyorganosiloxanes
- 15 • polyorganosiloxanes having at least one cationic or potentially cationic function in the medium (MAV)
- amphoteric polyorganosiloxanes having at least one cationic or potentially cationic function in the medium (MAV) and at least one function which is
20 neutral in the medium (MAV) and potentially anionic in the rinsing medium (MR)
- polyorganosiloxanes having at least one function which is neutral in the medium (MAV) and potentially anionic in the rinsing medium (MR).

25 7. Formulation according to any one of claims 1) to 6), characterized in that the material constituting the active substance (A) is an α - ω -

bis(hydroxy)polydimethylsiloxane, an α - ω -bis-(trimethyl)polydimethylsiloxane, a polymethylphenylsiloxane or a cyclic polydimethylsiloxane, preferably in oil form.

5 8. Formulation according to any one of claims 1) to 6), characterized in that the material constituting the active substance (A) is an amino polyorganosiloxane.

 9. Formulation according to claim 8),
10 characterized in that the amino polyorganosiloxane is a polyorganosiloxane which has hindered piperidyl groups.

 10. Formulation according to any one of claims 6) to 9), characterized in that said polyorganosiloxane is linear.

15 11. Formulation according to any one of claims 1) to 5), characterized in that the active substance (A) comprises a material selected from

- mono-, di- or triglycerides of C_1 - C_{30} carboxylic acids or mixtures thereof, such as vegetable oils
- 20 - sugar esters, sucroglycerides
- C_1 - C_{30} alcohol esters of C_1 - C_{30} carboxylic or C_2 - C_{30} dicarboxylic acids
- ethylene or propylene glycol monoesters or diesters of C_1 - C_{30} carboxylic acids
- 25 - propylene glycol C_4 - C_{20} alkyl ethers
- di(C_8 - C_{30} alkyl) ethers

- organic waxes comprising alkyl chains containing 4 to 40 carbon atoms.

12. Formulation according to any one of claims 1) to 11), characterized in that the ratio of
5 the mass of polymer constituting the active substance (A) to the mass of surfactant (TAC) is from 0.01 to 10, preferably from 0.01 to 1.

13. Formulation according to any one of claims 1) to 12), characterized in that the cationic
10 charges generated by the optional cationic or potentially cationic units of the material constituting the active substance (A) and by the cationic surfactant or surfactants at the surface of the material constituting the active active substance (A) in
15 dispersion in the medium (MAV) are such that the zeta potential of said polymer or copolymer in dispersion in (MAV) is from 0 to +50 mV, preferably from +10 to +40 mV.

14. Formulation according to any one of
20 claims 1) to 13), characterized in that the dispersion medium (MAV) for the active substance (A) is water or an aqueous-alcoholic polar medium.

15. Formulation according to claim 14),
characterized in that the alcohol or alcohols present
25 in the aqueous-alcoholic polar medium represent up to 70% of the volume of the medium (MAV).

16. Formulation according to any one of claims 1) to 15), characterized in that the polymer constituting the vehicle (V) is any polymer which is soluble or dispersible in aqueous or aqueous-alcoholic medium with a pH of between 2 and 8 and which comprises at least one unit which is neutral in the medium (MAV) and potentially anionic (HA) in the rinsing medium (MR).

17. Formulation according to claim 16), characterized in that the vehicle (V) polymer further comprises at least one unit which is cationic or potentially cationic (HC) in the medium (MAV) and/or at least one hydrophilic or hydrophobic nonionic unit.

18. Formulation according to any one of claims 1) to 17), characterized in that the relative amounts of the various units of the polymer constituting the vehicle (V) are such that in the medium (MAV) the overall charge of the polymer or copolymer is zero or cationic.

19. Formulation according to any one of claims 1) to 18), characterized in that the relative amounts of vehicle (V) polymer, surfactant (TAC), and material constituting the active substance (A) are such that in the course of the rinsing operation the number of anionic charges developed in the rinsing medium (MR) by the vehicle polymer (V) is sufficient to destabilize the active substance (A) in the rinsing medium (MR), in

particular by electrostatic attraction with the surface charges of the active substance (A) in the medium (MR).

20. Formulation according to claim 19), characterized in that the number of anionic charges developed in the rinsing medium (MR) by the vehicle (V) polymer to destabilize the active substance is at least 1% relative to the number of cationic surface charges of the active substance (A) in the medium (MR), and not more than 200% relative to the number of cationic surface charges of the active substance (A) in the medium (MR).

21. Formulation according to any one of claims 1) to 20), characterized in that the polymer constituting the vehicle (V) is a polymer selected from polymers derived from ethylenically unsaturated monomers, potentially anionic natural polysaccharides, potentially anionic or amphoteric substituted or modified polysaccharides, or mixtures thereof.

22. Formulation according to any one of claims 1) to 21), characterized in that the polymer constituting the vehicle (V) is a polymer derived:

- from at least one α - β monoethylenically unsaturated monomer which is neutral in the medium (MAV) and potentially anionic (HA) in the rinsing medium (MR) and

- optionally at least one α - β monoethylenically unsaturated monomer which is cationic or potentially cationic (HC) in the medium (MAV), and
- optionally at least one nonionic α - β monoethylenically unsaturated monomer which is hydrophilic or hydrophobic, preferably hydrophilic.

23. Formulation according to any one of claims 1) to 22), characterized in that the polymer constituting the vehicle (V) is a random, block or graft copolymer derived:

- from at least one α - β monoethylenically unsaturated hydrophilic monomer which is neutral in the medium (MAV) and potentially anionic (HA) in the rinsing medium (MR) and
- from at least one α - β monoethylenically unsaturated hydrophilic monomer which is cationic or potentially cationic (HC) in the medium (MAV),
- and optionally from at least one nonionic α - β monoethylenically unsaturated monomer which is hydrophilic or hydrophobic, preferably hydrophilic.

24. Formulation according to any one of claims 1) to 23), characterized in that the polymer constituting the vehicle (V) derives from one or more α - β monoethylenically unsaturated monomers and has an

average molar mass of greater than 5 000 g/mol,
preferably from 20 000 to 500 000 g/mol.

25. Formulation according to any one of
claims 1) to 24), characterized in that the polymer
5 constituting the vehicle (V) is selected from

- polyacrylic or polymethacrylic acids, alkali
metal polyacrylates or polymethacrylates,
preferably with a molar mass by weight of
from 100 000 to 1 000 000 g/mol
- 10 • acrylic acid/DADMAC copolymers, with a molar
ratio of 50/50 to 30/70, preferably with a
molar mass by weight of from 70 000 to
350 000 g/mol
- acrylic acid/MAPTAC copolymers, with a molar
15 ratio of 60/40 to 30/70, preferably with a
molar mass by weight of from 90 000 to
300 000 g/mol
- acrylic acid/MAPTAC/linear C₄-C₁₈ alkyl
methacrylate terpolymers comprising 0.005 to
20 10% by mass of alkyl methacrylate, with an
acrylic acid/MAPTAC molar ratio ranging from
60/40 to 30/70, and preferably having a molar
mass by weight of from 50 000 to
250 000 g/mol
- 25 • acrylic acid/dimethylaminoethyl methacrylate
(DMAEMA) copolymers, with a molar ratio of

60/40 to 30/70, preferably with a molar mass by weight of from 50 000 to 300 000 g/mol.

26. Formulation according to any one of claims 1) to 21), characterized in that the polymer
5 constituting the vehicle (V) is a potentially anionic natural polysaccharide formed of nonionic monosaccharide units and of monosaccharide units which are neutral in the medium (MAV) and potentially anionic in the rinsing medium (MR), and are alike or different.

10 27. Formulation according to claim 26), characterized in that said potentially anionic natural polysaccharide is a branched polysaccharide formed

- of a main chain comprising alike or different anhydrohexose units
- 15 • and of branches comprising at least one anhydropentose and/or anhydrohexose unit which is neutral in the medium (MAV) and optionally potentially anionic in the rinsing medium (MR).

20 28. Formulation according to claim 26) or 27), characterized in that said potentially anionic natural polysaccharide is a xanthan gum, a succinoglycan, a rhamsan, a gellan gum or a welan gum.

25 29. Formulation according to any one of claims 26) to 28), characterized in that said potentially anionic natural polysaccharide has a molar mass by weight of from 2 000 to 5 000 000, preferably

from 10 000 to 5 000 000, more particularly from 10 000 to 4 000 000 g/mol.

30. Formulation according to any one of claims 1) to 21), characterized in that the polymer
5 constituting the vehicle (V) is a substituted or modified polysaccharide whose native skeleton is formed of nonionic monosaccharide units and/or of monosaccharide units which are neutral in the medium (MAV) and potentially anionic in the rinsing medium
10 (MR), said monosaccharide units being alike or different and being substituted or modified

- by one or more groups which carry at least one charge which is neutral in the medium (MAV) and potentially anionic in the medium
15 (MR)
- and optionally by one or more groups which carry at least one charge which is cationic or potentially cationic in the medium (MAV), the degree of substitution or modification of the
20 monosaccharide units by the entirety of the groups which carry charges which are potentially anionic and of optional groups which carry cationic charges being such that said substituted or modified polysaccharide is soluble or dispersible
25 in aqueous or aqueous-alcoholic medium and has an overall cationic or zero charge in the medium (MAV).

31. Formulation according to claim 30), characterized in that said substituted or modified polysaccharide further comprises at least one nonionic modifying or substituent group.

5 32. Formulation according to claim 30) or 31), characterized in that said substituted or modified polysaccharide is a substituted or modified branched polysaccharide whose native skeleton is formed

- 10 • of a main chain comprising alike or different anhydrohexose units
- and of branches comprising at least one anhydropentose and/or anhydrohexose unit which is neutral in the medium (MAV) and optionally potentially anionic in the rinsing
- 15 medium (MR),

the anhydrohexose and/or anhydropentose units of said polysaccharide being substituted or modified by one or more groups which carry at least one charge which is neutral in the medium (MAV) and

20 potentially anionic in the medium (MR) and optionally at least one charge which is cationic or potentially cationic in the medium (MAV),

the degree of substitution or modification DS_i of the anhydrohexose and/or anhydropentose units by

25 the entirety of said groups which carry charges which are ionic or potentially ionic ranging from 0.01 to less than 3, preferably from 0.01 to 2.5,

with a ratio of the number of potentially anionic charges in the medium (MR) to the number of cationic or potentially cationic charges in the medium (MAV) ranging from 100/0 to 30/70, preferably from 100/0 to 50/50.

33. Formulation according to any one of claims 29) to 32), characterized in that said substituted or modified polysaccharide has a molar mass by weight of from 2 000 to 5 000 000, preferably from 10 000 to 5 000 000 g/mol.

34. Formulation according to any one of claims 29) to 33), characterized in that the native skeleton of said substituted or modified polysaccharide is a galactomannan.

35. Formulation according to any one of claims 29) to 34), characterized in that the native skeleton of said substituted or modified polysaccharide is selected from

- carboxymethylgalactomannans, especially carboxymethylguars,
- carboxymethylhydroxypropylgalactomannans, especially carboxymethylhydroxypropylguars,
- carboxymethyl-hydroxypropyltrimethylammonium chloride galactomannans, especially carboxymethyl-hydroxypropyltrimethylammonium chloride guars,
- carboxymethylhydroxypropyl-hydroxypropyl-trimethylammonium chloride galactomannans,

especially carboxymethyl-hydroxypropyl-
hydroxypropyltrimethylammonium chloride guar.

36. Formulation according to any one of
claims 1) to 35), characterized in that the amount of
5 vehicle (V) present in said formulation is from 0.01 to
5 parts by weight, preferably from 0.001 to 4 parts by
weight, and more particularly from 0.05 to 2 parts by
weight per 100 parts by weight of active substance (A).

37. Formulation according to any one of
10 claims 1) to 36), characterized in that it is in the
form of an aqueous or aqueous-alcoholic dispersion
comprising per 100 parts of its weight:

- from 0.01 to 40, preferably from 0.05 to 30
parts by dry weight of active substance (A)
- 15 - from 0.01 to 50, preferably from 0.01 to 35
parts by dry weight of surfactant (TAC)
- from 0.001 to 4, preferably from 0.01 to 1
part by dry weight of vehicle (V) polymer.

38. Formulation according to any one of
20 claims 1) to 37), characterized in that it further
comprises one or more customary constituents of
cationic rinsing formulations, selected from cationic
softeners, optical brighteners, color transfer
inhibitors, water-soluble monovalent mineral salts,
25 dyes, fragrances, foam suppressants, enzymes and
bleaches.

39. Process for treating textile fiber
articles by contacting said articles in the course of a
rinsing operation in aqueous or aqueous-alcoholic
medium with the rinsing formulation (F) of any one of
5 claims 1) to 38), and recovering said rinsed articles.

40. Process intended to enhance the
antiwrinkle and/or easy-iron and/or soil release and/or
abrasion resistance properties of textile fiber
articles, which consists in contacting said articles in
10 the course of a rinsing operation in aqueous or
aqueous-alcoholic medium with the rinsing formulation
(F) of any one of claims 1) to 38), and in recovering
said rinsed articles.

41. Use in a formulation (F) intended for
15 use in an operation of rinsing (R) textile fiber
articles (S) by means of an aqueous or aqueous-
alcoholic medium (MR), formulation (F) comprising at
least one active substance (A) comprising at least one
liquid or solid organic or organosilicon material in
20 particulate form and being

- in the form of a stable dispersion with a pH
of from 2 to 5 of said active substance (A) in an
aqueous or aqueous-alcoholic medium (MAV) or
- in a solid form obtained by drying said
25 dispersion,

the nature of the active substance (A) and of the aqueous or aqueous-alcoholic medium (MAV) being such that the active substance (A)

- is insoluble in the medium (MAV)
- 5 • has an overall zero or cationic charge in the medium (MAV),
- is stabilized in the medium (MAV) by means of a cationic surfactant (TAC), it being possible for said cationic surfactant (TAC)
- 10 to be wholly or partly replaced by a nonionic surfactant when the material constituting the active substance (A) is intrinsically cationic or intrinsically potentially cationic in the medium (MAV)
- 15 • remains insoluble in the rinsing medium (MR);

of at least one organic polymer which

- ♦ is soluble or dispersible in the medium (MAV) and in the rinsing medium (MR)
- 20 ♦ has an overall cationic or zero ionic charge in the medium (MAV)
- ♦ and is capable, at the pH of the rinsing operation in the rinsing medium (MR), of developing anionic charges in sufficient
- 25 quantity to destabilize the active substance (A) in the rinsing medium (MR);

as a vehicle (V) capable of bringing said active substance (A) toward the surface of said textile fiber articles (S) in the rinsing operation (R).

42. Process for enhancing the deposition of
5 an active substance (A) comprising at least one solid or liquid organic or organosilicon material in particulate form on the surface of textile fiber articles (S), during an operation of rinsing of said articles by means of an aqueous or aqueous-alcoholic
10 medium (MR) obtained from a formulation (F) comprising said active substance (A), the formulation (F) being
- in the form of a stable dispersion with a pH of from 2 to 5 of said active substance (A) in an aqueous or aqueous-alcoholic medium
15 (MAV) or
 - in a solid form obtained by drying said dispersion,
- the nature of the active substance (A) and of the aqueous or aqueous-alcoholic medium (MAV) being such that the active substance (A)
- 20
- is insoluble in the medium (MAV)
 - has an overall zero or cationic charge in the medium (MAV),
 - is stabilized in the medium (MAV) by means of
25 a cationic surfactant (TAC), it being possible for said cationic surfactant (TAC) to be wholly or partly replaced by a nonionic

surfactant when the material constituting the active substance (A) is intrinsically cationic or intrinsically potentially cationic in the medium (MAV)

- 5 • remains insoluble in the rinsing medium (MR);

by adding to said formulation (F) a vehicle (V)

comprising at least one organic polymer which

- 10 ♦ is soluble or dispersible in the medium (MAV)
 and in the rinsing medium (MR)
- ♦ has an overall cationic or zero ionic charge
 in the medium (MAV)
- ♦ and is capable, at the pH of the rinsing
 operation in the rinsing medium (MR), of
15 developing anionic charges in sufficient
 quantity to destabilize the active substance
 (A) in the rinsing medium (MR).

20 43. Processes according to any one of claims
 40), 41) or 43), or use according to claim 42),
 characterized in that the amount of formulation
 employed, expressed in terms of dry matter, is
 from 0.001 to 5 g/l, preferably from 0.05 to 2 g/l
 in the rinsing bath.